

98

99.

5

10

A method of immunizing against *Neisseria* comprising administering to a host an immunizing amount of an immunogenic composition comprising a polypeptide having an amino acid sequence of SEQ ID NO:2 or a fragment thereof, a polypeptide having an amino acid sequence of SEQ ID NO:4 or a fragment thereof, a polypeptide having an amino acid sequence of SEQ ID NO:14 or a fragment thereof, a polypeptide having an amino acid sequence of SEQ ID NO:16 or a fragment thereof, a polypeptide having an amino acid sequence of SEQ ID NO:25 or a fragment thereof and a pharmaceutically acceptable carrier, wherein the amino acid at residue 18 of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:14, SEQ ID NO:16 and SEQ ID NO:25 is an amino acid other than an ATC encoded isoleucine.

99

100.

The method of claim 99, wherein the amino acid at residue 18 is tyrosine.

15

101.

100

The method according to any one of claims 44-53b, further comprising an adjuvant.

101

102.

20

The method according to any one of claims 44-53b, further comprising one or more PorA polypeptides or proteins selected from the group consisting of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18 and SEQ ID NO:20.

102

103.

25

An immunogenic composition according to any one of claims 42, 67, 76 or 85, further comprising one or more ORF2086 protein antigens comprising an amino acid sequence of SEQ ID NO:26 through SEQ ID NO:83.